



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Determinants of non-adherence to anti-tuberculosis treatment in high income, low incidence, settings**

**Citation for published version:**

on behalf of the IMPACT study group , Jones, ASK, Bidad, N, Horne, R, Stagg, HR, Wurie, FB, Kielmann, K, Karat, A, Kunst, H, Campbell, CNJ, Darvell, M, Clarke, AL & Lipman, MCI 2021, 'Determinants of non-adherence to anti-tuberculosis treatment in high income, low incidence, settings: A scoping review', *International Journal of Tuberculosis and Lung Disease*. <https://doi.org/10.5588/ijtld.21.0024>

**Digital Object Identifier (DOI):**

[10.5588/ijtld.21.0024](https://doi.org/10.5588/ijtld.21.0024)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

International Journal of Tuberculosis and Lung Disease

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



**Determinants of non-adherence to anti-TB treatment in high income, low TB incidence settings: a scoping review**

Annie S. K. Jones<sup>1</sup>, Natalie Bidad<sup>1</sup>, Rob Horne<sup>1</sup>, Helen R. Stagg<sup>2</sup>, Fatima B. Wurie<sup>3,4</sup>, Karina Kielmann<sup>5</sup>, Aaron S. Karat<sup>5, 6</sup>, Heinke Kunst<sup>7</sup>, Colin N. J. Campbell<sup>8</sup>, Marcia Darvell<sup>8</sup>, Amy L. Clarke,<sup>1</sup> Marc C. I. Lipman<sup>8,9</sup>, on behalf of the IMPACT study group (NIHR 16/88/06)

<sup>1</sup>Centre for Behavioural Medicine, Research Department of Practice and Policy, UCL School of Pharmacy, London, UK (ORCID ID: 0000-0001-7868-2804).

<sup>2</sup>Usher Institute, University of Edinburgh, Edinburgh, UK

<sup>3</sup>Research Department of Epidemiology and Public Health, Institute of Epidemiology and Health Care, UCL, London, UK

<sup>4</sup>Migrant Health, Public Health England, London, UK

<sup>5</sup>Institute for Global Health and Development, Queen Margaret University, Edinburgh, UK

<sup>6</sup>TB Centre, London School of Hygiene & Tropical Medicine, London, UK

<sup>7</sup>Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK

<sup>8</sup>UCL Respiratory, Division of Medicine, UCL, London, UK

<sup>9</sup>Royal Free London NHS Foundation Trust, London, UK

**IMPACT study group:**

Professor Ibrahim Abubakar

Professor Andrew Copas

Mr Mike Mandelbaum

Dr Alistair Story

Dr Caroline Clark

Corresponding Author: Professor Marc C. I. Lipman, UCL Respiratory, Division of Medicine, UCL, London, UK. [marclipman@nhs.net](mailto:marclipman@nhs.net)

Running head: Determinants of anti-TB treatment adherence

Key words: tuberculosis, adherence, treatment, determinants

35 Word count: 2,499

36

## **ABSTRACT**

### *Background*

Improving adherence to anti-tuberculosis (TB) treatment is a public health priority in high income, low incidence (HILI) regions. We conducted a scoping review to identify reported determinants of non-adherence in HILI settings.

### *Methods*

Key terms related to tuberculosis, treatment, and adherence were used to search MEDLINE, EMBASE, Web of Science, PsycINFO, and CINAHL in June 2019. Quantitative studies examining determinants (demographic, clinical, health systems, or psychosocial) of non-adherence to anti-TB treatment in HILI settings were included.

### *Results*

From 10,801 results, we identified 24 relevant studies from 10 countries. Definitions and methods of assessing adherence were highly variable, as were documented levels of non-adherence (0.9%–89%). Demographic factors were assessed in all studies and clinical factors frequently assessed (23/24). Determinants commonly associated with non-adherence were homelessness, imprisonment, and alcohol or drug misuse. Health system (8/24) and psychosocial factors (6/24) were less commonly evaluated.

### *Conclusion*

Our review identified some key factors associated with non-adherence to anti-TB treatment in HILI settings. Modifiable determinants such as psychosocial factors are under-evidenced and should be further explored as these may be better targeted by adherence support. There is an urgent need to standardise definitions and measurement of adherence to more accurately identify the strongest determinants.

## INTRODUCTION

Despite the availability of effective, low-cost medication, tuberculosis (TB) remains a global health concern <sup>1</sup>. One reason for this is non-adherence to anti-TB treatment, which increases morbidity and mortality <sup>2,3</sup>, transmission, the development of drug resistance, and health disparity <sup>4–6</sup>.

We have yet to identify the best adherence support for anti-TB treatment. Directly-observed therapy (DOT) has been recommended by the World Health Organization (WHO) since the 1990s <sup>7</sup>, but research does not consistently find DOT superior to self-administered therapy (SAT) in reducing adverse treatment outcomes such as loss to follow-up <sup>8,9</sup>. Furthermore, improved outcomes from DOT dissipate when patients receiving SAT have increased contact with healthcare services <sup>8</sup>, suggesting the benefit of DOT may result from the “encounter” rather than the “observation”. This is important as DOT is resource-intensive, and can be perceived negatively by patients <sup>10–12</sup>.

Interventions to support adherence are more likely to be effective if they address the specific causes of non-adherence relevant to the individual patient <sup>13,14</sup>. Identifying specific, and potentially modifiable, determinants of adherence to anti-TB treatment is therefore critical in developing more targeted and effective support <sup>15</sup>.

Improving anti-TB treatment adherence is a priority for high income, low TB incidence (HILI) countries progressing toward TB elimination <sup>16</sup>. To date, determinants have mostly been examined in high incidence regions <sup>17–19</sup>. Determinants in high and low incidence regions may differ, based on differences in populations with TB and resources for care <sup>20–22</sup>. Therefore, as formative research for an intervention to promote adherence to anti-TB treatment in the UK <sup>23</sup>, we undertook a scoping review to explore determinants of non-adherence to anti-TB treatment within HILI settings, and identify evidence gaps relevant to patients and healthcare providers to be addressed by future research.

## METHODS

We selected a scoping review methodology to provide a broad overview and highlight key evidence gaps<sup>24</sup>, given expectations of study heterogeneity<sup>25,26</sup> and diverse definitions and measurements of TB treatment adherence<sup>27</sup>. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension Checklist for Scoping Reviews (PRISMA-ScR) was used<sup>25</sup>.

### *Literature search*

Five databases (MEDLINE, EMBASE, Web of Science, PsycINFO, and CINAHL) were searched in June 2019. Researchers developed and refined search terms related to TB, treatment, and adherence, with support from an experienced librarian (Supplementary Material 1).

Search terms were mapped to the Population-Concept-Context framework recommended for scoping reviews<sup>28</sup> (Table 1). Identified studies were imported into Endnote<sup>29</sup> and duplicates were removed. Two authors independently screened titles and abstracts using the website Rayyan, designed for article screening in reviews<sup>30</sup>. Any discrepancies were resolved through discussion. Reference lists of included studies were hand-searched to identify additional relevant studies.

Eligibility criteria are listed in Table 1. Included studies were peer-reviewed, English language studies, whose aim was to report primary, observational, quantitative data on determinants of non-adherence to anti-TB treatment, in countries classified as high income<sup>31</sup> with low TB incidence rates (<40 per 100,000 people), when the study was conducted. We included outcomes of both discontinuation (early cessation of treatment, including loss to follow-up) and suboptimal implementation (missing doses during treatment)<sup>32–34</sup>. We excluded qualitative research, as our research group has reviewed this separately<sup>35</sup>.

### *Data extraction and synthesis*

Two authors independently extracted data (cross-checking 50% of studies). Determinants were included if studied as primary exposures of interest or potentially confounding factors. Determinants were labelled as demographic, clinical, health systems-related, or psychosocial.

Categorises were used to reflect the strength of evidence for each determinant. A proxy measure was created for this, based on the size and direction of the effect size (ES) estimate and statistical certainty. Evidence was classified from strongest to weakest using the following categories:

- Category 1: **Strongest:** ES (ratio)  $\geq 1.5$ , p-value  $\leq 0.05$ ;
- Category 2: ES (ratio)  $\geq 1.5$ , p-value  $> 0.05$ , small sample size ( $n < 154$ ) i.e. study likely to be under-powered;
- Category 3: ES (ratio)  $> 1.0$  to  $< 1.5$ , p-value  $\leq 0.05$ ;
- Category 4: **Weakest:** ES (ratio)  $> 1.0$ , p-value  $> 0.05$ .

The equivalent categories were used to classify determinants observed to have a protective effect. In order to provide a standardised classification for category 2, a sample size calculation was required. It was calculated that a minimum of 154 participants would indicate an adequately powered sample size, using 90% power and 5% significance level, statistically conservatively assuming that 50% of individuals had the outcome among the unexposed, and assuming a one-to-one ratio of exposed to unexposed or cases to controls. Although this threshold did not perfectly reflect the analyses in all studies, it provided a framework for weighting the evidence of each determinant. It did not indicate judgement on the quality of included studies. Where possible, determinants were classified based on ES in multivariable, not univariable, analyses.

## *Ethics*

Ethics approval was not required as this was a scoping review.

## **RESULTS**

### *Description of included studies*

The initial search found 10,801 studies. After removing duplicates, 9,932 remained for title and abstract screening, and 25 met the inclusion criteria (Figure 1, Supplementary Material 2). Data on determinants were extracted for 24 studies, as one<sup>36</sup> reported no ES.

Included studies were published 1986-2019, from 10 different countries, including the UK and Ireland ( $n=7$ )<sup>37-43</sup>, USA ( $n=6$ )<sup>36,44-48</sup>, and Spain ( $n=5$ )<sup>49-53</sup>. The most

common study design was retrospective cohort (n=12)<sup>37,40,46,48,49,53–59</sup>. Sample sizes ranged from 62 to 73,591 (median= 1009; interquartile range (IQR)= 184-2576). The mean/median participant age ranged from 28.0 to 52.1 years. The median percentage of males was 64.4% (IQR= 56.0-71.0%).

Most studies (n=20) included all patients starting treatment in a given setting<sup>36–45,47,48,50–52,54–58</sup>. Three studies sampled specific high-risk groups, of people experiencing homelessness or unstable living arrangements<sup>60</sup>, individuals with multidrug-resistant TB (MDR-TB)<sup>59</sup>, or HIV/TB co-infection<sup>53</sup>. Two studies compared outcomes between groups within a cohort, such as immigrants versus individuals born within a country<sup>46,49</sup>.

#### *Non-adherence: definitions and assessment*

Supplementary Material 2 demonstrates the considerable variability in definitions of adherence. Most study outcomes (n=15) related to treatment discontinuation (stopping treatment early)<sup>38,40,44–47,49,50,53–56,58–60</sup>. Fewer study outcomes (n=7) appeared to record suboptimal implementation (missed doses during treatment)<sup>36,37,39,41,43,48,57</sup>. One study included both a discontinuation and suboptimal implementation outcome<sup>42</sup>. Two studies used a single outcome encapsulating both discontinuation and suboptimal implementation<sup>51,52</sup>.

Discontinuation outcomes were often measured using state or national registries/surveillance databases<sup>40,44–47,53–55,58,59</sup>, hospital/lab records<sup>44,45,47,49,59</sup>, or medical notes<sup>38,60</sup>.

Sub-optimal implementation was assessed using various methods, including adherence scale scores<sup>36</sup>, medical records<sup>57</sup>, physician impression from interviews/assessments<sup>39,41</sup>, patient self-report<sup>39</sup>, health visitor reports (including pill counts)<sup>41</sup>, urine samples (to detect rifampicin)<sup>39,43</sup>, attendance at appointments<sup>41,48,57</sup>, and prescription requests<sup>48,57</sup>.

Overall, retrospective studies most often used surveillance/registry data to determine adherence<sup>37,40,45–47,53–55,58,59</sup>, whereas prospective studies used more varied



methods (Supplementary Material 2). Reported non-adherence ranged from 0.9% to 89% across studies (median= 7.0%; IQR= 5.2-16.3%). Two studies did not report levels of non-adherence <sup>36,60</sup>.

### *Determinants of non-adherence*

#### *Demographic determinants*

Demographic determinants were assessed by all 24 studies (Supplementary Material 2). Specifically, the most studied determinant groups were place of residence and age (Supplementary Material 3). The variable with the greatest strength of evidence for a large effect on non-adherence (Categories 1 or 2- large effect sizes with  $p\text{-value}\leq 0.05$  or a small sample size, see Methods; Supplementary Material 2) was place of residence (Figure 2). Within that variable, homelessness <sup>37,42,44,46–48,50,53,57,60</sup> and living in an institution or prison (e.g. a “confined institution”, a residence hall, or mental hospital) <sup>37,42,46,47,51,52,55,58</sup> had the strongest evidence, weighted overall, towards non-adherence (Supplementary Material 2).

Age, sex, ethnicity, and nationality also showed mixed evidence of effects, as within each variable just as many or more studies found a weak effect with non-adherence as a large effect (Figure 2). Ethnicity and nationality determinants appeared very context-specific, demonstrated by the variation in baseline comparators within these categories. Overall, few demographic determinants were classified in categories 2 (i.e. large ES,  $p\text{-value}>0.05$ , but small sample size) or 3 (small ES,  $p\text{-value}\leq 0.05$ ) in terms of strength of evidence. The grouping variables most commonly found to have a weak effect on adherence (category 4, small ES,  $p\text{-value}>0.05$ ) were age, nationality/origin, and ethnicity. No variable had a consistently large effect with non-adherence.

#### *Clinical determinants*

Clinical determinants were the second most studied category (23/24 studies Supplementary Material 2). The substance use/misuse grouping variable was the most frequently assessed and had the most evidence weighted towards a large effect (Supplementary Material 3 and Figure 2). Specifically, illicit drug misuse/addiction had the strongest evidence for this <sup>48,50–52,55</sup> (Supplementary Material 2). The evidence for clinical determinants was also mixed, in terms of both

strength of evidence and direction. For example, in the HIV grouping variable, HIV positive status was a risk factor for non-adherence<sup>50,51,55</sup>, yet a diagnosis of AIDS was protective against non-adherence<sup>44,46</sup> (Supplementary Material 4). Again, few determinants fell into categories 2 and 3 in terms of strength of evidence, and the grouping variables which most commonly showed a weak effect with adherence were smear and sputum result, substance use/misuse, and HIV infection.

#### *Health systems determinants*

Health systems determinants were less frequently investigated (8/24 studies). Within this category, route to care was the most studied grouping variable (Supplementary Material 3). Healthcare professionals' perception of patient understanding (e.g. lack of awareness of TB severity, understanding of treatment instructions, language barriers) had a consistently large effect with non-adherence, though this determinant was minimally studied. The grouping variables most often found to have a weak effect with adherence were route to care, and those classified as "other".

#### *Psychosocial determinants*

Psychosocial determinants were the least studied (6/46 studies), where only mental health and having close relationships were assessed (Supplementary Material 3, Figure 2). Of these, having a mental health problem was both a risk for<sup>42</sup> and protective against non-adherence<sup>47</sup> (Supplementary Material 2). Strength of evidence for mental health problems was also mixed, with as many studies finding strong and weak effects on adherence.

## **DISCUSSION**

In our scoping review investigating the determinants of non-adherence to anti-TB treatment within HILI settings, homelessness, imprisonment, and alcohol or drug misuse were commonly associated factors. Health systems and psychosocial determinants were under-explored. Considerable heterogeneity in measurements and definitions of non-adherence was present across studies, hindering the conclusions that can be drawn.

When synthesising the literature on determinants, we found that demographic and clinical factors were most studied. This may reflect the relative ease of capturing this

data through TB surveillance in HILI settings, such as the UK <sup>61</sup>. However, the context required to understand mixed findings for these determinants was largely missing from studies, which may result from utilising these data sources. Without context, these findings are unhelpful for explaining non-adherence. For example, a recent systematic review found that despite assumptions, non-adherence was as likely to occur in both migrants and non-migrants <sup>62</sup>. Such findings highlight the importance of contextualising demographic and clinical determinants, if researchers are to utilise this data in intervention design.

In addition, demographic and clinical determinants are largely non-modifiable (e.g. history of imprisonment) or difficult to change (such as homelessness, illicit drug use/addiction) within a feasible, scalable healthcare intervention <sup>63</sup>. Improving adherence to anti-TB treatment requires identifying potentially modifiable determinants that can be targeted within a pragmatic, person-centred healthcare intervention.

Determinants more amenable to change, such as health systems issues, have rarely been quantitatively assessed in HILI settings. Health systems barriers in high incidence regions, such as distance to treatment facilities and transport costs <sup>17–19</sup>, may be less apparent in HILI countries with better-resourced health services. Nonetheless, they may affect subgroups of patients, given that TB disproportionately affects people with lower socioeconomic status in high income settings <sup>20</sup>. In addition, health systems determinants may interact with other factors (such as fear of stigma making an individual seek care at a more distant hospital), reinforcing the need to better understand their influence in HILI settings.

Psychosocial determinants are also under-researched in quantitative literature on TB adherence. This oversight is significant given the known relationship between TB, stigma, and adherence, even in low incidence settings <sup>64</sup>. Understanding the social context of TB treatment is significant for reaching TB control goals, given the well-established links between social determinants of health and inequality <sup>65</sup>, even within regions of low TB incidence <sup>66</sup>.

Theory in behavioural medicine suggests adherence is best viewed as a modifiable behaviour and not a trait <sup>67</sup>, as adherence patterns can change within an individual over time <sup>32,68</sup>, and also differ between people with shared demographic characteristics. Theory and evidence suggest that amendable, cognitive and affective factors, such as beliefs about illness and treatment, influence subsequent coping strategies, including treatment adherence <sup>69–71</sup>. Understanding psychosocial determinants may enable us, therefore, to provide better adherence support.

Evidence from this review has important clinical implications for intervention development in TB. Interventions should: 1) accurately assess known risk factors for non-adherence to anti-TB treatment in HILI settings; and 2) mitigate the influence of these on perceptual and practical barriers to adherence <sup>70</sup>. For example, interventions should be tailored to both target a patient's beliefs about TB and treatment, and provide practical support to overcome personal barriers to treatment.

Our scoping review followed PRISMA-ScR guidelines to systematically search the available literature. We may have been limited by including only English language studies. We may have missed secondary data reported (e.g. in intervention studies) by only including studies whose primary aim was to examine determinants of non-adherence. In addition, as this was a scoping review, the quality of included studies was not assessed.

Our understanding of non-adherence to anti-TB treatment within HILI settings is severely limited by the heterogeneity of included studies. Clearer and consistent definitions of which type of non-adherence is being assessed in studies <sup>33</sup>, and data presented beyond simple binary summary measures, are urgently needed <sup>72</sup>.

By including all data on reported determinants, whether measured as primary exposures of interest or potential confounding factors, some estimates may be subject to bias. Of note, few (n=2) studies assessed all four categories of determinants and therefore adjusted for confounders appropriately. This considerably impairs our ability to understand the interaction between determinants and their relationship to non-adherence, and may explain the inconsistency of the included evidence.

334

335 In conclusion, this scoping review identifies determinants with the best supportive  
336 evidence, and highlights a gap in our understanding of adherence to anti-TB  
337 treatment in HILI settings. Understanding how demographic and clinical  
338 determinants are associated with adherence to anti-TB treatment is necessary to  
339 inform intervention development. Qualitative work could extend current  
340 understanding by examining how health systems and psychosocial factors influence  
341 anti-TB treatment in HILI settings <sup>23</sup>. Stakeholders in TB policy and service  
342 implementation should also consider how factors influencing patient adherence are  
343 currently evaluated and understood. Existing care practices, such as risk  
344 assessments, should ensure the range of complex factors involved in adherence are  
345 comprehensively addressed.

346

347 We also identified a need for greater consistency in definitions and measurement of  
348 adherence within the TB literature. Without this, it will remain difficult to effectively  
349 synthesise data, and understand reported patterns of adherence behaviour.

350

## ACKNOWLEDGEMENTS

ASK reports grants from the National Institute for Health Research (UK), during the conduct of the study; personal fees from The Aurum Institute (South Africa), The Center for Health Policies and Studies (Republic of Moldova), Edanz Group (Japan), Vital Strategies (Singapore), University of Cape Town (South Africa), the Bill & Melinda Gates Foundation (USA), and Bloomberg Philanthropies (USA) outside the submitted work; non-financial support from Kyoto University (Japan) and the Africa Health Research Institute (South Africa) outside the submitted work. CNJC reports personal fees from Public Health England outside the submitted work. HRS reports grants from Medical Research Council (UK) and the National Institute for Health Research (UK) during the conduct of the study; other from Korean CDC and Johnson and Johnson and other from Latvian Society Against Tuberculosis outside the submitted work; and HRS is a core group member of the World Health Organization's European Tuberculosis Research Initiative and co-chair of UK Academics and Professionals Against Tuberculosis. KK, MCIL, and MD report grants from the National Institute for Health Research (NIHR) during the conduct of the study. RH is supported by the National Institute for Health Research (NIHR, Collaboration for Leadership in Applied Health Research and Care, North Thames at Bart's Health NHS Trust and Asthma UK (AUKCAR). Speaker engagements with honoraria with the following companies: Abbvie, Amgen, Astellas, AstraZeneca, Biogen, Erasmus, Idec, Gilead Sciences, GlaxoSmithKline, Janssen, Merck Sharp Dohme, Novartis, Pfizer, Roche, Shire Pharmaceuticals, and TEVA. RH is founding director of a UCL-Business spin-out company (Spoonful of Sugar Ltd) providing consultancy on treatment engagement and patient support programmes to healthcare policy makers, providers, and industry. FBW is an employee of Public Health England. All other authors declare no conflicts of interest.

This work was supported by the National Institute for Health Research (NIHR) Health Technology Assessment Programme, UK grant number 16/88/06. The views expressed are those of the authors and not necessarily those of the National Health Service, UK, the National Institute for Health Research or the Department of Health and Social Care. HRS is funded by the Medical Research Council (MRC), UK [MR/R008345/1].

AJ, NB, RH, HRS, FBW and MCIL conceived of and designed the work. All authors acquired, analysed, or interpreted the data. AJ drafted the paper. All authors revised it critically for important intellectual content. All authors give final approval of the version to be published and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity are appropriately investigated and resolved.

## REFERENCES

1. United Nations. High-level Meeting on the Fight Against Tuberculosis [Internet]. New York, NY.; 2018. Available from: <https://www.who.int/tb/unhlmonTBDeclaration.pdf>
2. Nahid P, Jarlsberg LG, Rudoy I, de Jong BC, Unger A, Kawamura LM, et al. Factors associated with mortality in patients with drug-susceptible pulmonary tuberculosis. *BMC Infect Dis.* 2011;11(1):1.
3. Zerbini E, Greco A, Estrada S, Cisneros M, Colombo C, Beltrame S, et al. Risk factors associated with tuberculosis mortality in adults in six provinces of Argentina. *Medicina (B Aires).* 2017;77(4):267–73.
4. Pradipta IS, Forsman LD, Bruchfeld J, Hak E, Alffenaar JW. Risk factors of multidrug-resistant tuberculosis: A global systematic review and meta-analysis. *J Infect.* 2018;77(6):469–78.
5. Rockwood N, Abdullahi LH, Wilkinson RJ, Meintjes G. Risk factors for acquired rifamycin and isoniazid resistance: A systematic review and meta-analysis. *PLoS One.* 2015;10(9):1–23.
6. Hayward S, Harding RM, McShane H, Tanner R. Factors influencing the higher incidence of tuberculosis among migrants and ethnic minorities in the UK. *F1000Research.* 2018;7.
7. World Health Organization. Communicable Diseases Cluster. What is DOTS ? A Guide to Understanding the WHO-recommended TB Control Strategy Known as DOTS. World Health Organization; 1999.
8. Karumbi J, Garner P. Directly observed therapy for treating tuberculosis. *Cochrane Database Syst Rev.* 2015;2015(5).
9. Van De Berg S, Jansen-Aaldring N, De Vries G, Van Den Hof S. Patient support for tuberculosis patients in low-incidence countries: A systematic

review. PLoS One. 2018;13(10):1–24.

10. Hansel NN, Wu AW, Chang B, Diette GB. Quality of life in tuberculosis: Patient and provider perspectives. *Qual Life Res.* 2004;13(3):639–52.
11. Sagbakken M, Bjune GA, Frich JC. Humiliation or care? A qualitative study of patients' and health professionals' experiences with tuberculosis treatment in Norway. *Scand J Caring Sci.* 2012;26(2):313–23.
12. Craig GM, Zumla A. The social context of tuberculosis treatment in urban risk groups in the United Kingdom: A qualitative interview study. *Int J Infect Dis.* 2015;32:105–10.
13. Horne R, Weinman J, Barber N, Elliot R, Morgan M, Cribb A. Concordance, adherence and compliance in medicine taking: Report for the National Coordinating Centre for NHS Service Delivery and Organisation R & D (NCCSDO). London, UK; 2005.
14. National Institute for Health and Care Excellence. Medicines adherence: involving patients in decisions about prescribed medicines and supporting adherence Clinical guideline. 2009.
15. Pradipta IS, Houtsma D, van Boven JFM, Alffenaar J-WC, Hak E. Interventions to improve medication adherence in tuberculosis patients: a systematic review of randomized controlled studies. *npj Prim Care Respir Med.* 2020;30(1):21.
16. Lönnroth K, Migliori GB, Abubakar I, D'Ambrosio L, De Vries G, Diel R, et al. Towards tuberculosis elimination: An action framework for low-incidence countries. *Eur Respir J.* 2015;45(4):928–52.
17. Castelnovo B. A review of compliance to anti tuberculosis treatment and risk factors for defaulting treatment in Sub Saharan Africa. *Afr Health Sci.* 2010;10(4):320–4.
18. Tola HH, Tol A, Shojaeizadeh D, Garmaroudi G. Tuberculosis treatment non-adherence and lost to follow up among TB patients with or without HIV in developing countries: A systematic review. *Iran J Public Health.* 2015;44(1):1–11.
19. Zegeye A, Dessie G, Wagnaw F, Gebrie A, Mohammed S, Islam S, et al. Prevalence and determinants of anti-tuberculosis treatment non-adherence in Ethiopia: A systematic review and meta-analysis. *PLoS One.* 2019;14(1):e0210422.
20. Public Health England. Tuberculosis in England: 2019 report. Public Health



- England, London; 2019.
21. Centers for Disease Control and Prevention. Reported Tuberculosis in the United States, 2018 [Internet]. 2019. Available from: <https://www.cdc.gov/tb/statistics/reports/2018/default.htm>
22. Zumla A, Raviglione M, Hafner R, Von Reyn CF. Tuberculosis. *N Engl J Med*. 2013;368(8):745–55.
23. Stagg HR, Abubakar I, Campbell CNJ, Copas A, Darvell M, Horne R, et al. IMPACT study on intervening with a manualised package to achieve treatment adherence in people with tuberculosis: Protocol paper for a mixed-methods study, including a pilot randomised controlled trial. *BMJ Open*. 2019;9(12):e032760.
24. Peterson J, Pearce PF, Ferguson LA, Langford CA. Understanding scoping reviews: Definition, purpose, and process. *J Am Assoc Nurse Pract*. 2017;29(1):12–6.
25. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Ann Intern Med*. 2018;169(7):467–73.
26. Tricco AC, Lillie E, Zarin W, O'Brien K, Colquhoun H, Kastner M, et al. A scoping review on the conduct and reporting of scoping reviews. *BMC Med Res Methodol*. 2016;16(15).
27. DiMatteo MR. Variations in patients' adherence to medical recommendations: A quantitative review of 50 years of research. *Med Care*. 2004;42(3):200–9.
28. Peters M, Godfrey C, McInerney P, Soares CB, Khalil H, Parker D. Methodology for JBI scoping reviews. In: *The Joanna Briggs Institute Reviewers Manual 2015*. South Australia: The Joanna Briggs Institute; 2015. p. 3–24.
29. EndNote 7.0. Thomson Reuters; 2015.
30. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev*. 2016;5(1).
31. World Bank Group. World Bank Country and Lending Groups [Internet]. 2020. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>
32. Stagg HR, Lewis JJ, Liu X, Huan S, Jiang S, Chin DP, et al. Temporal Factors

- and Missed Doses of Tuberculosis Treatment. A Causal Associations Approach to Analyses of Digital Adherence Data. *Ann Am Thorac Soc*. 2020;17(4):438–49.
33. Stagg HR, Flook M, Martinecz A, Kielmann K, Wiesch PA Zur, Karat AS, et al. All non-adherence is equal, but is some more equal than others? TB in the digital era. *ERJ Open Res*. 2020;
  34. World Health Organization. Definitions and reporting framework for tuberculosis-2013 revision. 2013.
  35. Arakelyan S, Karat AS, Vidal N, Stagg HR, Jones ASK, Darvell M, et al. Relational dynamics of anti-tuberculosis treatment-related behaviour in high-income countries: a scoping review. *Submiss*.
  36. McDonnell M, Turner J, Weaver MT. Antecedents of adherence to antituberculosis therapy. *Public Health Nurs*. 2001;18(6):392–400.
  37. Anderson C, Anderson SR, Maguire H, Hayward AC, Story A. Tuberculosis in London: The convergence of clinical and social complexity. *Eur Respir J*. 2016;48(4):1233–6.
  38. Anyama N, Bracebridge S, Black C, Niggebrugge A, Griffin SJ. What happens to people diagnosed with tuberculosis? A population-based cohort. *Epidemiol Infect*. 2007;135(7):1069–76.
  39. Corcoran R. Compliance with chemotherapy for tuberculosis. *Ir Med J*. 1986;79(4):87–90.
  40. Millett ERC, Noel D, Mangtani P, Abubakar I, Kruijsaar ME. Factors associated with being lost to follow-up before completing tuberculosis treatment: Analysis of surveillance data. *Epidemiol Infect*. 2013;141(6):1223–31.
  41. Ormerod LP, Prescott RJ. Inter-relations between relapses, drug regimens and compliance with treatment in tuberculosis. *Respir Med*. 1991;85(3):239–42.
  42. Story A, Murad S, Roberts W, Verheyen M, Hayward AC. Tuberculosis in London: The importance of homelessness. problem drug use and prison. *Thorax*. 2007;62(8):667–71.
  43. Wardman AG, Knox AJ, Muers MF, Page RL, Leeds Chest Clinic. Profiles of non-compliance with therapy. *Br J Dis Chest*. 1988;82(3):285–9.
  44. Brudney K, Dobkin J. Resurgent Tuberculosis in New York City : Human Immunodeficiency Virus , Homelessness , and the Decline of Tuberculosis

- Control Programs. *J Public Health Policy*. 1992;13(4):435–50.
45. Cook S V., Fujiwara PI, Frieden TR. Rates and risk factors for discontinuation of rifampicin. *Int J Tuberc Lung Dis*. 2000;4(2):118–22.
46. Cummings KC, Mohle-Boetani J, Royce SE, Chin DP. Movement of tuberculosis patients and the failure to complete antituberculosis treatment. *Pneumologie*. 1998;52(11):625–6.
47. Driver CR, Matus SP, Bayuga S, Winters AI, Munsiff SS. Factors associated with tuberculosis treatment interruption in new york city. *J Public Heal Manag Pract*. 2005;11(4):361–8.
48. Pablos-Méndez A, Knirsch CA, Barr RG, Lerner BH, Frieden TR. Nonadherence in tuberculosis treatment: Predictors and consequences in New York City. *Am J Med*. 1997;102(2):164–70.
49. Ballesteros AL, Oriol J, Francisco I, Fernández S, García Bragado F, Vinyes A. Clinical characteristics of tuberculosis in immigrants and autochthonous populations in two hospitals of Catalonia. *Rev Clínica Española*. 2014;214(8):445–52.
50. Caylà JA, Caminero JA, Rey R, Lara N, Vallés X, Galdós-Tangüis H. Current status of treatment completion and fatality among tuberculosis patients in Spain. *Int J Tuberc Lung Dis*. 2004;8(4):458–64.
51. Caylà JA, Rodrigo T, Ruiz-Manzano J, Caminero JA, Vidal R, García JM, et al. Tuberculosis treatment adherence and fatality in Spain. *Respir Res*. 2009;10:1–11.
52. Rodrigo T, Caylà JA, Casals M, García-García JM, Caminero JA, Ruiz-Manzano J, et al. A predictive scoring instrument for tuberculosis lost to follow-up outcome. *Respir Res*. 2012;13:1–9.
53. Galdós Tangüis H, Caylà JA, García De Olalla P, Jansà JM, Brugal MT. Factors predicting non-completion of tuberculosis treatment among HIV-infected patients in Barcelona (1987-1996). *Int J Tuberc Lung Dis*. 2000;4(1):55–60.
54. Kawatsu L, Uchimura K, Ohkado A, Kato S. A combination of quantitative and qualitative methods in investigating risk factors for lost to follow-up for tuberculosis treatment in Japan – Are physicians and nurses at a particular risk? *PLoS One*. 2018;13(6):1–13.
55. Nunes C, Duarte R, Veiga AM, Taylor B. Who are the patients that default

- tuberculosis treatment? - Space matters! *Epidemiol Infect.* 2017;145(6):1130–4.
56. Tetart M, Meybeck A, Assaf A, Valette M, Choisy P, Blondiaux N, et al. Factors of loss to follow-up during tuberculosis treatment in a low-incidence region. *Med Mal Infect.* 2020;50(1):28–35.
  57. Zellweger JP, Coulon P. Outcome of patients treated for tuberculosis in Vaud County, Switzerland. *Int J Tuberc Lung Dis.* 1998;2(5):372–7.
  58. Borgdorff MW, Veen J, Kalisvaart NA, Broekmans JF, Nagelkerke NJD. Defaulting from tuberculosis treatment in the Netherlands: Rates, risk factors and trend in the period 1993-1997. *Eur Respir J.* 2000;16(2):209–13.
  59. Jensenius M, Winje BA, Blomberg B, Mengshoel AT, Von Der Lippe B, Hannula R, et al. Multidrug-resistant tuberculosis in Norway: A nationwide study, 1995-2014. *Int J Tuberc Lung Dis.* 2016;20(6):786–92.
  60. Kizuki M, Takano T, Nakamura K, Fukuda Y, Watanabe M, Inose T, et al. Social Course Patterns of Urban Dwellers with Tuberculosis under Fragile Living Conditions in Tokyo, Japan. *J Epidemiol.* 2006;16(4):1–5.
  61. Royal College of Nursing. A Case Management Tool for TB Prevention, Care and Control in the UK. Royal College of Nursing, London, UK; 2019.
  62. Nellums LB, Rustage K, Hargreaves S, Friedland JS. Multidrug-resistant tuberculosis treatment adherence in migrants: A systematic review and meta-analysis. *BMC Med.* 2018 Feb 22;16(1):27.
  63. Sabaté E. Adherence to long-term therapies: Evidence for action. Geneva, Switzerland: World Health Organization; 2003.
  64. Craig GM, Daftary A, Engel N, O'Driscoll S, Ioannaki A. Tuberculosis stigma as a social determinant of health: a systematic mapping review of research in low incidence countries. Vol. 56, *International Journal of Infectious Diseases.* Elsevier B.V.; 2017. p. 90–100.
  65. Marmot M. Social determinants of health inequalities. *Lancet.* 2005 Mar 19;365(9464):1099–104.
  66. Marmot M, Allen J, Bell R, Bloomer E, Goldblatt P. WHO European review of social determinants of health and the health divide. Vol. 380, *The Lancet.* Lancet Publishing Group; 2012. p. 1011–29.
  67. Chan A, Horne R. Beliefs and Adherence in Hypertension and Cardiovascular Protection. In: *Drug Adherence in Hypertension and Cardiovascular Protection.*

- 589 Springer, Cham; 2018. p. 123–41.
- 590 68. Kielmann K, Vidal N, Riekstina V, Krutikov M, Werf MJV, Biraua E, et al.  
591 “treatment is of primary importance, and social assistance is secondary”: A  
592 qualitative study on the organisation of tuberculosis (TB) care and patients’  
593 experience of starting and staying on TB treatment in Riga, Latvia. PLoS One.  
594 2018;13(10).
- 595 69. Leventhal H, Meyer D, Nerenz D. The common sense representation of illness  
596 danger. In: Rachman S, editor. Contributions to medical psychology. New  
597 York: Pergamon Press; 1980. p. 7–30.
- 598 70. Horne R, Cooper V, Wileman V, Chan A. Supporting Adherence to Medicines  
599 for Long-Term Conditions: A Perceptions and Practicalities Approach Based  
600 on an Extended Common-Sense Model. Eur Psychol. 2019;24(1):82–96.
- 601 71. Foot H, La Caze A, Gujral G, Cottrell N. The necessity-concerns framework  
602 predicts adherence to medication in multiple illness conditions: A meta-  
603 analysis. Patient Educ Couns. 2016;99(5):706–17.
- 604 72. Vernon A, Fielding K, Savic R, Dodd L, Nahid P. The importance of adherence  
605 in tuberculosis treatment clinical trials and its relevance in explanatory and  
606 pragmatic trials. PLOS Med. 2019 Dec 10;16(12):e1002884.  
607

608 Table 1. Patient-Concept-Context (PCC) elements and inclusion/exclusion criteria.

PCC element	Definition in scoping review	Inclusion criteria	Exclusion criteria
Population	Patients taking anti-tuberculosis treatment	Studies reporting data on non-adherence to treatment for pulmonary tuberculosis.	<ul style="list-style-type: none"> <li>• Studies with a non-human sample.</li> <li>• Studies with patients taking prophylactic TB treatment or treatment for latent TB.</li> <li>• Studies where the majority of patients (&gt;50%) had extra-pulmonary disease.</li> <li>• Studies with a co-morbid sample (excluding HIV).</li> </ul>
Concept	Determinants of non-adherence to treatment	Peer-reviewed studies reporting primary, observational, data on determinants of non-adherence to treatment.	<ul style="list-style-type: none"> <li>• Studies reporting interventions (including studies where DOT/VOT were standard treatment, or more than 50% of the sample was receiving DOT/VOT).</li> <li>• Qualitative studies.</li> <li>• Studies that were not primary research articles (e.g. reviews, commentaries, or letters).</li> <li>• Studies that did not measure determinants of non-adherence.</li> <li>• Studies where treatment completion was the outcome (as this is conflated with successful treatment outcome and is not a measure of patient adherence).</li> </ul>
Context	HILI TB settings	Countries classified as high income and low TB incidence at time of study.	<ul style="list-style-type: none"> <li>• Studies in settings defined as low and middle income, or with high TB incidence.</li> </ul>


609 Note. DOT= directly-observed therapy; HILI= high income, low (TB) incidence; TB=  
610 tuberculosis; VOT= video-observed therapy.  
611

## List of Figures

Figure 1. PRISMA diagram of screening process and included studies.


Figure 2. Determinants of non-adherence to TB treatment. Note. Bars may include multiple determinant levels assessed within the same study. Darkest grey indicates the strongest effect (i.e. category 1: a large risk or protective effect at  $p < .05$ ), medium grey indicates a large risk or protective effect at  $p > .05$  with a small sample size (category 2), light grey indicates a small risk or protective effect at  $p < .05$  (category 3), and lightest grey indicates the weakest effect found at  $p > .05$  (category 4). HCP = healthcare professional, SES= socioeconomic status, TB= tuberculosis

**Figure 1.**





650 **Figure 2**  
651



652

**Determinants of non-adherence to anti-tuberculosis treatment in high income,  
low incidence, settings: A scoping review**  
**Annie S. K. Jones, Natalie Bidad, Rob Horne Helen R. Stagg, Fatima B. Wurie,  
Karina Kielmann, Aaron S. Karat, Heinke Kunst, Colin N. J. Campbell, Marcia  
Darvell, and Marc Lipman, on behalf of the IMPACT study group (NIHR  
16/88/06)**

**Online Supplement**

## Supplementary material 1

### Scoping review search example strategy from MEDLINE

1. Tuberculosis/
2. (TB or tuberculo\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, organism supplementary concept word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
3. 1 or 2
4. Drug Therapy/
5. (medication\* or medicine\* or treatment\* or therap\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, organism supplementary concept word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
6. 4 or 5
7. 3 and 6
8. Antitubercular Agents/
9. 7 or 8
10. Medication Adherence/ or "Treatment Adherence and Compliance"/
11. (adheren\* or complian\* or non-adheren\* or non-complian\* or nonadheren\* or concordan\* or non-concordan\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, organism supplementary concept word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
12. (LTFU or "los\* to follow-up" or "los\* to follow up" or LFU or default).mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, organism supplementary concept word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
13. Lost to Follow-Up/
14. 10 or 11 or 12 or 13
15. 9 and 14
16. limit 15 to (english language and humans)

## Supplementary material 2. Bibliometric data from included studies (n=25).

Year(s) of study	Sample size (N)	Country	Study population	Study design	Categories of determinants assessed	Adherence measure	Non-adherence definition	% Non-adherent	Citation information
<b>Prospective designs</b>									
Not given	90	Ireland	Patients being treated for pulmonary TB and discharged from a Dublin hospital.	Prospective cohort	Demographic Clinical	Urine sample, self-report or physician's impression from interview	Non-compliance: based on interview evidence and $\geq 1$ negative urine sample(s).	23.3	Corcoran <sup>39</sup>
Not given	113	UK	Patients being treated at Leeds Chest Clinic receiving a rifampicin-containing regimen.	Prospective cohort	Demographic Clinical	Urine sample	Non-compliance: negative urine sample.	7.0	Wardman et al. <sup>43</sup>
1988-1989	224	USA	Patients being treated in Harlem Hospital Center, New York.	Prospective cohort	Demographic Clinical	State TB registry data and hospital records	Noncompliance: no follow-up treatment or LTFU.	89.0	Brudney & Dobkin <sup>44</sup>
1995-1996	62	USA	Patients being treated and residing within Georgia.	Prospective cohort	Psychosocial	Tuberculosis General Adherence Scale (TBGAS)	Lower scores on TBGAS scale	Not given (mean TBGAS score= 92.6%)	McDonnell et al. <sup>36*</sup>
1999-2000	1515	Spain	Patients being treated by a member of the Tuberculosis and Respiratory Infections Group of the Sociedad Española de Neumología y Cirugía Torácica (SEPAR).	Prospective cohort	Demographic Clinical Health systems	Epidemiological questionnaire completed by staff, including assessments of "appointment attendance, physician estimation, and patient confirmation" (data source not specified)	Default: no treatment received for >1 month or missed appointments.	4.0	Cayla et al. <sup>50</sup>

Year(s) of study	Sample size (N)	Country	Study population	Study design	Categories of determinants assessed	Adherence measure	Non-adherence definition	% Non-adherent	Citation information
1998–2003	119	Japan	Homeless patients or those in fragile living situations who received treatment in a Tokyo hospital (excluded patients with HIV/TB co-infection).	Prospective cohort	Demographic Clinical Health systems Psychosocial	Medical notes	Treatment interruption during outpatient care: no treatment for ≥2 consecutive months.	Not given (19.33 worked out from results)	Kizuki et al. <sup>60</sup>
2000–2003	575	UK	Patients being treated in the East of England.	Prospective cohort	Demographic Clinical	Patient notes (extracted by TB staff)	LTFU	7.8	Anyama et al. <sup>38</sup>
2003	1941	UK	Patients in Greater London who were or should have been on treatment.	Prospective cohort	Demographic Clinical Psychosocial	Self-report, pill counts, urine tests, medical records, case-manager “knowledge” of patient	Poor adherence: self-reported, inconsistent pill counts, negative urine test, or patients switched to DOT or hospitalised for poor adherence.  LTFU: being out of contact with services for at ≥2 months without medication during first 6 months of treatment.	Poor adherence: 46.0  LTFU: 15.0	Story et al. <sup>42</sup>
2006–2007	1490	Spain	Patients being treated by a member of the Tuberculosis and Respiratory Infections Group of the Sociedad Española de Neumología y Cirugía Torácica (SEPAR) (excluded patients with known drug resistance or those not initiating standard treatment).	Prospective cohort	Demographic Clinical Health systems	Electronic diary completed by staff (no details regarding from where data obtained)	Poor adherence: including default (treatment interruption for >2 months, non-completion by 9-months on standard regimen, or <80% prescribed doses taken) and LTFU.	6.2	Cayla et al. <sup>51</sup>

Year(s) of study	Sample size (N)	Country	Study population	Study design	Categories of determinants assessed	Adherence measure	Non-adherence definition	% Non-adherent	Citation information
2006–2009	1490	Spain	Patients being treated with culture-positive or smear-positive disease, extrapulmonary TB with caseating granuloma, identification by histology, or clinical, radiological, epidemiological or laboratory suspicion of TB (excluded patients with known drug-resistance or those with a contraindication to start standard treatment).	Prospective cohort	Demographic Clinical Health systems	Not specified	LFTU: treatment interruption (any reason) for $\geq 2$ months, non-completion of treatment within 9 months for standard therapy, or taking $< 80\%$ of prescribed dose.	6.48	Rodrigo et al. <sup>52</sup>
<b>Retrospective designs</b>									
1988–1992	103	Switzerland	Patients with bacteriologically confirmed pulmonary TB being treated in Vaud County.	Retrospective cohort	Demographic Clinical Psychosocial	Questionnaire completed by practitioners (using medical records)	Not adherent: not specified (adherence considered satisfactory if patient attended scheduled visits and requested prescriptions).	18.4	Zellweger & Coulon <sup>57</sup>
1993	2576	USA	Compared patients being treated in California who did and did not move during treatment to another health jurisdiction.	Retrospective cohort	Demographic Clinical Psychosocial	National TB surveillance data	Default: patients who refused treatment or were LTFU.	5.5	Cummings et al. <sup>46</sup>
1991–1994	184	USA	Patients with a first time, positive-culture being treated in New York City.	Retrospective cohort	Demographic Clinical	Contacting providers for clinic attendance and prescription information	Noncompliance: not attending clinic appointments for $\geq 2$ months, or $\geq 3$ months during 1 year.	48.0	Pablos-Mendez et al. <sup>48</sup>
1993–1994	3520	USA	Patients with culture-confirmed, rifampin- susceptible TB, starting a rifampin-containing regimen of at least 60 days, being treated in New York City.	Retrospective case-control	Demographic Clinical Health systems	State TB registry data and hospital records	Inappropriate treatment discontinuation: discontinuing rifampicin without experiencing serious adverse effects related to use.	0.9	Cook et al. <sup>45</sup>

Year(s) of study	Sample size (N)	Country	Study population	Study design	Categories of determinants assessed	Adherence measure	Non-adherence definition	% Non-adherent	Citation information
1987–1996	1354	Spain	Patients with HIV/TB co-infection, detected by the Active Epidemiological Surveillance System of the Barcelona Tuberculosis Prevention and Control Programme, being treated in Barcelona.	Retrospective cohort	Demographic Clinical	National TB surveillance data	Treatment abandonment: LTFU or failed medical controls and not found by public health surveillance nursing team.	13.1	Galdós Tangüis et al. <sup>53</sup>
1993–1997	7529	The Netherlands	Patients being treated in the Netherlands.	Retrospective cohort	Demographic Clinical Health systems	National TB registry data	LTFU (excludes patients reportedly continuing treatment elsewhere)	8	Borgdorff et al. <sup>58</sup>
1998–2002	328	USA	Patients who were culture-positive being treated in New York City (excluded patients with MDR-TB).	Retrospective case-control	Demographic Clinical Health systems Psychosocial	State TB registry data, patient interview forms, hospital records (including case manager notes)	Default (treatment interrupted for ≥60 days) with return to therapy  Default without return to therapy (including LTFU or treatment refusal)	4.2	Driver et al. <sup>47</sup>
2001–2007	41,120	UK	Patients being treated in England, Wales and Northern Ireland reported to the Enhanced Tuberculosis Surveillance (ETS) system.	Retrospective cohort	Demographic Clinical	National TB surveillance data	LTFU (before treatment completion, including patients who moved overseas)	5.9	Millet et al. <sup>40</sup>
2000–2011	503	Spain	Compares immigrant and native patients being treated in Catalonia.	Retrospective cohort	Demographic Clinical Psychosocial	Hospital records	Treatment abandonment: treatment interrupted for ≥2 months (without medical advice), or LTFU with no information available.	1.8	Ballesteros et al. <sup>49</sup>
2009–2012	12,908	UK	Patients being treated in London, England.	Retrospective cohort	Demographic Clinical	National TB surveillance and laboratory data (matched with national outreach data - “find and treat” registry)	Non-adherence (not specified)	5.6	Anderson et al. <sup>37</sup>

Year(s) of study	Sample size (N)	Country	Study population	Study design	Categories of determinants assessed	Adherence measure	Non-adherence definition	% Non-adherent	Citation information
2000–2013	27,894	Portugal	Patients with pulmonary TB being treated in continental Portugal, as identified through the national TB surveillance database (SVIG-TB).	Retrospective cohort	Demographic Clinical	National TB surveillance data	Default: treatment interrupted for >8 weeks after completing ≥1 month of treatment)	4.9	Nunes et al. <sup>55</sup>
1995–2014	68	Norway	Patients with MDR-TB being treated in Norway.	Retrospective cohort	Demographic Clinical	Hospital and laboratory records, TB registry data	LTFU: WHO 2013 definition (no treatment initiation, or treatment interrupted for ≥2 consecutive months).	17.6	Jensenius et al. <sup>59</sup>
2006–2015	73,591	Japan	Patients with pulmonary TB being treated in Japan.	Retrospective cohort	Demographic Clinical Health systems	National TB surveillance data	LTFU: definition from Japanese TB surveillance system (treatment interrupted for ≥2 consecutive months, or treatment duration <6 months).	7.8	Kawatsu et al. <sup>54</sup>
1997–2017	190	France	Patients diagnosed at Dron Hospital in Tourcoing (excluding those with MDR-TB or XDR-TB).	Retrospective cohort	Demographic Clinical	Not specified (appears to be medical and laboratory records)	LTFU: no treatment initiation or treatment interrupted for ≥2 consecutive months.	15.0	Tetart et al. <sup>56</sup>
<b>Mixed designs</b>									
1978–1987	1009	UK	Patients being treated in Blackburn, England.	Retrospective and prospective cohort	Demographic	Physician assessment, monthly health visitor reports (including pill counts) and clinic attendance	Poor compliance: ≥3 missed appointments or unfavourable assessments.	3.0	Ormerod & Prescott <sup>41</sup>

**Note.** \*determinants not extracted for this study. TB= Tuberculosis; LTFU= Loss to follow-up; MDR-TB= Multidrug-resistant tuberculosis; WHO = World Health Organization; XDR-TB= Extensively drug-resistant tuberculosis.



### Supplementary material 3. Frequency of determinants assessed by included studies.

Determinant category	Determinant grouping variable	Studies assessing determinant grouping variable:
Demographic	Age	n= 14 37 38 40 41 42 46 47 49 50 51 53 58 59 55
	Sex	n= 14 38 40 42 43 44 48 50 51 53 57 58 59 54 55
	Ethnicity	n= 5 40 42 45 46 48
	Nationality/origin	n= 7 37 38 43 48 58 59 54
	Residency/Immigration status	n= 9 40 49 50 51 52 56 57 58 55
	Place of residence (e.g. homelessness, history of living in an institution or prison)	n= 16 37 42 44 46 47 48 49 50 51 52 53 57 58 54 55 60
	Employment status	n= 4 51 56 54 60
	Occupation	n= 2 46 54

<b>Determinant category</b>	<b>Determinant grouping variable</b>	<b>Studies assessing determinant grouping variable:</b>
	SES indices (e.g. primary education only, living in low SES neighbourhood)	n= 5 38 39 43 53 54
	Other (e.g. relationship status, receiving leave for appointments, place of residency, travelling behaviour, moved health jurisdiction during treatment)	n= 4 39 40 46 59
Clinical	Disease site/presentation	n= 6 37 38 40 48 56 54
	Smear & sputum culture result	n= 6 38 47 48 57 58 54
	Treatment resistance	n= 3 46 48 60
	TB history	n= 9 37 38 40 43 45 52 53 56 60
	HIV status	n= 9 44 46 48 49 50 51 53 56 55
	Substance use/issues	n= 17 37 39 42 44 45 46 47 48 50 51 52 53 56 57 59 55 60
	Other (e.g. relapse (unspecified, diabetes co-morbidity)	n= 2 57 60

<b>Determinant category</b>	<b>Determinant grouping variable</b>	<b>Studies assessing determinant grouping variable:</b>
Health systems	Route to care	n= 3 51 58 60
	Treatment provider (e.g. treated at referral hospital, treated by private health provider)	n= 2 45 54
	HCP perceptions of patient knowledge	n=3 47 51 52
	Other (e.g. hospitalisation, health insurance status, time from culture confirmation to presentation)	n= 3 47 50 54
Psychosocial factors	Mental health problems	n= 3 42 47 57
	Having close relationships	n= 1 60

Note. HCP= health care professional, SES= socioeconomic status, TB= tuberculosis

## Supplementary material 4

### Strength of evidence for demographic, clinical, health-systems and psychosocial factors associated with adherence to TB treatment

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR $\geq$ 1.5, P $\leq$ 0.05	OR/RR $\geq$ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P $\leq$ 0.05	OR/RR< 1.5, P> 0.05		OR/RR $\leq$ 0.5, P $\leq$ 0.05	OR/RR $\leq$ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P $\leq$ 0.05	OR/RR> 0.5, P> 0.05
<b>Demographic Factors</b>										
Age	Under 30 y/o	37			42§,					
		<u>[16-24]</u>			<u>[30-59]</u>					
		51*								
		<u>[&gt;50]</u>								
		59								
	25-34 y/o	<u>[<math>\geq</math>40]</u>								
		41*								
		[30-44, 45-59, >60]								
					38*					
					<u>[0-24]</u>					
30-65 y/o	30-65 y/o			58¶	51*				40	37
				[<25]	<u>[&gt;50]</u>				<u>[15-44]</u>	<u>[16-24]</u>
					53*				55	38*
					<u>[15-29]</u>				<u>[15-34]</u>	<u>[0-24]</u>
					58***					
	Over or equal 65 y/o				<u>[&lt;25]</u>					
					42§,					
					<u>[30-59]</u>					
							37		40	38*
							<u>[16-24]</u>		<u>[15-44]</u>	<u>[0-24]</u>
Other (e.g. unspecified, broad range)	Other (e.g. unspecified, broad range)	59		50*	53*					
		<u>[<math>\geq</math>40]</u>		[ $\leq$ 17]	<u>[15-29]</u>				<u>[15-34]</u>	38*
		49			58				<u>[0-24]</u>	47***
		<u>[&lt;40]</u>							[NS]	
							<u>[16-24]</u>		<u>[15-44]</u>	
	Other (e.g. unspecified, broad range)						<u>[15-29]</u>		<u>[15-34]</u>	
							50*		55	
							<u>[15-29]</u>		<u>[15-34]</u>	
							<u>[15-29]</u>		<u>[15-34]</u>	
							<u>[15-29]</u>		<u>[15-34]</u>	

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05		OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
		46††,††			[<25]					[NS]
		[>45]								
		58§§								
		[<25]								
Sex	Male [Female]	50*	59*	40	44	Male [Female]				44
		51*		58	48*					43*
		55*			53*					54
		42§,			38*					
		57*								
Ethnicity	Hispanic	45*			48* [White]					
		[non-Hispanic Black]								
		46††,††								
	Asian [White]	[Asian]								
		40*,†††		40*,†††	48*	Asian [White]				40*,§§§
					42					42§
		46††,††			42§,    [White]					
	Black African/ Caribbean/Black British/non-Hispanic black	[Asian]								
		48* [White]								
		40* [White]								
	Non-Hispanic White	45*								
		[non-Hispanic Black]								
		45								
	Ethnicity category non-specific (e.g. White, Asian, or Hispanic) [NS]									
Nationality/ origin‡	Europe	40*			42	Other [White]				42§
						Black Caribbean [White]				42
		37		37¶¶¶¶	37****					
		[South Asia]		[South Asia]	[South Asia]					
		58								
		[Dutch]								

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05		OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
	Asia		43* [British]		37 [South Asia] 58 [Dutch]					
	North America and Oceania [Abroad]				48*	North America and Oceania [South Asia]				37
	Africa	58†††† [Dutch]	59 [NS]		37 [South Asia] 58†††† [Dutch]					
	East Mediterranean [Dutch]	58				East Mediterranean [South Asia]				37
	Foreign-born [UK born]	38*				Foreign-born [Japan born]				54
	Other/country of birth unknown [Dutch]	58				Other/Country of birth unknown [Japan born]		54		
	Latin, South, Central America or Caribbean				37 [South Asia] 58 [Dutch]					
Residency/ immigration status	Immigrant or migrant [native]	49				North Africa [South Asia]				37
		50				Immigrant or migrant [native]				56*
		51								
		55								
		52								
		57*								
	Recent migrant (under 4 years)	40*			58 [UK born] [other]					
	Migrant 5+ years [UK born]				40*					
	Illegal immigrant [not in category]	58								
	Asylum seeker [not in category]				58					
	Time in resident country unknown [UK born]	40*								

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05		OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
Place of residence	Living alone	51								
		<u>[with family]</u> 52								
		<u>[NS]</u>								
	Homelessness/no fixed abode <i>[has fixed abode]</i>	37	60		58	Homelessness/no fixed abode <i>[has fixed abode]</i>			55	54
		44	57*		49					
		50*								
		46††,††								
		47†††,***								
		60§§§§								
		48								
		53								
		42§,								
History/living in institution or prison	History/living in institution or prison	52		53*	50*					
		<u>[NS]</u> 58		<i>[no history]</i>	<u>[NS]</u> 46††					
		<u>[no history]</u> 55*			<i>[not in category]</i>					
		<u>[no history]</u> 47†††								
		<u>[not incarcerated]</u> 42§								
		<u>[no imprisonment during current treatment]</u> 37								
		<u>[NS]</u> 51								
		<u>[with family]</u>								

Grouping variable	Potential risk factor	Strength of evidence			
		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05
		46†††			
		[not in category]			
	Shared accommodation [with family]	51			
	Living in a county jail at diagnosis	46††,††	51 [with family]		
		[not in category]			
Employment status	Active occupational status [retired]	51*			
	Unemployed	51* [retired]	60* [employed]		
	Disabled occupational status [retired]	51*			
Occupation	HCP [Full/part- time employed]		54††††††	54*****	
	Housemaker [Full/part- time employed]			54	
	Job/employment unknown [Full/part- time employed]		54		
	Migrant agricultural work [NS]	46††,††			
SES indices	Receiving social welfare benefit [not in category]			54	
	Low SES level neighbourhood [any other SES level neighbourhood]	53			
	Townsend score high deprivation [Townsend score least deprivation]			38*	

Potential protective factor	Strength of evidence			
	1	2	3	4
	OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
Unemployed [Full/part- time employed]				54
Temporary/self employed	56* [NS]			54 [Full/part- time employed]
HCP [Full/part- time employed]				54†††††
Student [Full/part- time employed]				54



Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR $\geq$ 1.5, P $\leq$ 0.05	OR/RR $\geq$ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P $\leq$ 0.05	OR/RR< 1.5, P> 0.05		OR/RR $\leq$ 0.5, P $\leq$ 0.05	OR/RR $\leq$ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P $\leq$ 0.05	OR/RR> 0.5, P> 0.05
	Primary education only [2nd/3rd level education only]	39*								
	SES Level 4/5 (high deprivation) [SES Level 1,2, or 3]	43*								
	Having medical card [no medical card]	39*								
						Townsend score between low and high deprivation [Townsend score least deprivation]				38*
Other	Living outside of London [living in London]			40*						
	Living in Oslo [NS]		59							
	Single/separated or widowed [married]	39*								
	Moved health jurisdiction within state during Tx [not moving during Tx]	46*								
<b>Clinical Factors</b>										
Disease site/ presentation	Pulmonary [extra-pulmonary]	37		40		Pulmonary [extra-pulmonary]				38*
	Extra-pulmonary [not in category]	56*			48*					
						With cavity			54 [no cavity]	48* [no cavitory disease]
Smear & sputum culture	Smear +ve and/or culture +ve				38* [-ve]	Smear +ve and/or culture +ve	47*** [not +ve in first 30 days of initial sputum collection]		54 [-ve]	

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05		OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
					47¶¶¶ [not +ve in first 30 days of initial sputum collection] 57*		58 [no bacteriological confirmation]			
	Culture/smear unknown/not done			54††††† [-ve]	48* [+ve] 54§§§§§ [-ve]					
						Smear -ve and/or culture – ve [+ve]				48*
T <sub>x</sub> resistance	MDR	46††,†† [NS]			48* [No resistance]					
	Other resistance [no resistance]				48*	Other resistance [no resistance]				60*
TB history	Previous TB	37 [NS]	60* [no history]		40 [no history]	Previous TB [no history]				38*
	Previous TB T <sub>x</sub> [no previous T <sub>x</sub> ]	45 52 53 43*								
	Unknown previous TB status [no history]			40						
	Previous TB T <sub>x</sub> default [no previous default]	43*								
HIV status	HIV infection [HIV negative]	50* 51* 55			48*	First episode of TB [NS] HIV infection [HIV negative]	56			49*
	HIV status known/missing [HIV negative]				51* 48*					
	HIV negative [NS]	56								
	AIDS [HIV negative]				48*		44			

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05		OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
						AIDS (or AIDS related complex) <i>[NS]</i>	46††			46††
	HIV infection via IDU transmission <i>[sexual transmission]</i>	53*								
Substance use/misuse	Alcohol use <i>[NS]</i>	37								
	Alcohol misuse/addiction	44	39*		48*					
		<i>[NS]</i>		<i>[0 drinks per week]</i>		<i>[no history of alcoholism]</i>				
		55	57*		53*					
		<i>[not in category]</i>		<i>[NS]</i>		<i>[No alcoholism]</i>				
		56*								
		<i>[NS]</i>								
	Illicit drug use	37	57*		48*		Illicit drug use <i>[NS]</i>			47***
		<i>[NS]</i>	<i>[NS]</i>		<i>[No cocaine use]</i>					
		42			47¶¶¶					
		<i>[NS]</i>			<i>[NS]</i>					
		59								
	<i>[NS]</i>									
Illicit drug misuse/addiction	51									
	<i>[No IDU]</i>									
	55									
	<i>[NS]</i>									
	48									
	<i>[no IDU]</i>									
	52									
	<i>[NS]</i>									
	50									
	<i>[not drug addict]</i>									
Drug use unknown	51									
	<i>[No IDU]</i>									
	52									
	<i>[NS]</i>									

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05		OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
	Any substance misuse (including IDU, non-IDU, and alcohol) [NS]	46††,‡‡								
	Being treated with methadone [NS]	45								
	Alcohol problems in hospital [Not in category]		60*							
Other	Relapse (unspecified) [NS]	57*								
						Diabetes co-morbidity [not in category]				60*
<b>Health-Systems Factors</b>										
Route to care	Source – emergencies [primary care]	51*								
	Source – specialist [primary care]				51*					
	Source – other [primary care]				51*					
	Collapsing on street [other reason]		60*							
	Detection of TB by screening [other detection]				58					
Tx provider	T <sub>x</sub> started in OP department [initial hospitalisation]			54						
	Private health provider [provider was DOH]	45*								
	Private health provider with low volume of patients [private health provider with high volume]	45*								
	T <sub>x</sub> by low volume provider [NS]	45								
HCP perception of	Had previous T <sub>x</sub> comprehension† [no previous Tx comprehension]	51								

Grouping variable	Potential risk factor	Strength of evidence				Potential protective factor	Strength of evidence			
		1	2	3	4		1	2	3	4
		OR/RR≥ 1.5, P≤ 0.05	OR/RR≥ 1.5, P> 0.05, n< 154	OR/RR> 1.0 to <1.5, P≤ 0.05	OR/RR< 1.5, P> 0.05		OR/RR≤ 0.5, P≤ 0.05	OR/RR≤ 0.5, P> 0.05, n< 154	OR/RR> 0.5 to <1.0, P≤ 0.05	OR/RR> 0.5, P> 0.05
patient's knowledge	Had difficulty with previous Tx comprehension† <i>[easy previous Tx comprehension]</i>	51								
	Poor understanding <i>[NS]</i>	52								
	Lack of awareness of TB severity <i>[NS]</i>	47¶¶¶,***								
Other	Hospitalised (includes IP care) <i>[not hospitalised]</i>				50*					
	Months from +ve culture to DOH interview† <i>[NS]</i>			47¶¶¶		Months from +ve culture to DOH interview <i>[NS]</i>				47***
	No health insurance <i>[has health insurance]</i>				54					
<b>Psychosocial Factors</b>										
Mental health issues	Mental health problems <i>[NS]</i>	42§,				Mental health problems <i>[NS]</i>	47***			47¶¶¶ 57*
Having close relationships						Having close relationships <i>[no close relationships]</i>		60*		

**Note.** Where variable levels are non-binary, baseline comparator is given italicised in square brackets, either next to variable level or individual study reference where this differs between studies. No data was extracted from <sup>36</sup>. Some variables could not be extracted from <sup>58</sup> (urban residence, previous default from TB Tx, homelessness, alcohol addiction, drug addiction, occupation, travel to endemic areas, disease site, HIV co-infection), <sup>57</sup> (age), <sup>43</sup> (age, nationality (other)), <sup>60</sup> (sex, age, disease site, cavitory disease, sputum smear result), and <sup>39</sup> (drinking (moderate drinking)). +ve= positive, -ve= negative, DOH= Department of Health, HCP= healthcare professional, IDU= intravenous drug use, IP= inpatient, MDR= multidrug-resistant, NS=not specified, OP= outpatient, SES= socioeconomic status, TB= tuberculosis, Tx= Treatment \*=univariate/ bivariate analysis. †=Determinants were not further defined. ‡=nationality: studies <sup>37</sup>, <sup>48</sup>, and <sup>38</sup> comparator is not the study country, for studies <sup>58</sup>, <sup>59</sup>, <sup>43</sup> and <sup>54</sup>, comparator is study country. §=outcome: outcome: non-adherent in first 2 months, ||= outcome: loss to follow-up within 6 months, ¶= age: 35-44, \*\*= age: 55-64, ††= outcome: excludes patients who moved during study, ‡‡= outcome: includes patients who moved during study, §§= age: 75, |||= age: 45-54, ¶¶¶= outcome: default with return to therapy, \*\*\*= outcome: default without return to therapy, †††= ethnicity: Indian, ‡‡‡= ethnicity: Pakistani, §§§= ethnicity: Bangladeshi, ||||= nationality/origin: born in Central Europe, ¶¶¶¶= nationality/origin: born in West Europe, \*\*\*\*= nationality/origin: born in East Europe, ††††= nationality/origin: Somalian and other

African, ‡‡‡‡= nationality/origin: Moroccan, §§§§= place of residence: staying in transient hostel after discharge, ||||| = place of residence: homeless prior to admission, ¶¶¶¶= HCP: Nurse, \*\*\*\*\*= HCP: Physician, †††††= HCP: Other HCP, ‡‡‡‡‡= Smear: culture/smear not done, §§§§§= Smear: culture/smear unknown.